

MASSIVE STARS IN TRANSITION

Paul A. Crowther¹

Abstract. We discuss the various post-main sequence phases of massive stars, focusing on Wolf-Rayet stars, Luminous Blue Variables, plus connections with other early-type and late-type supergiants. End states for massive stars are also investigated, emphasising connections between Supernovae originating from core-collapse massive stars and Gamma Ray Bursts.

1 Introduction

Material introduced in my article dealing with OB stars is built upon here for post-main sequence phases of massive stars, namely Wolf-Rayet stars, slash stars, Luminous Blue Variables (LBVs), B[e] supergiants and cool supergiants. What are the properties and interrelations between these extreme objects? What happens to these stars at the end of their short lives?

The most dramatic explosions in the Universe, Gamma Ray Bursts (GRBs), convert a sizable fraction of a Solar Mass into energy within seconds. The currently favoured scenario for most GRBs is via the collapse of a massive star to form a black hole, with a corresponding Supernova explosion. Closer to home, η Car, the archetypal LBV is one of the most extreme Milky Way stars and briefly was the second brightest star in the night sky in the middle of the 19th century, despite lying 8000 light years from the Sun. During this eruption it ejected material at an *average* rate of one Earth mass every 15 minutes for two decades. Wolf-Rayet (W-R) stars, the evolved descendants of massive OB stars throw off several solar masses over lifetimes of 10^5 yr. W-R stars, plus their subsequent SN explosions are major contributors to the chemical enrichment of galaxies (Leitherer et al. 1992; Dray et al. 2003). It is apparent that the advanced stages of the most massive stars represent exceptional astrophysical objects, which are discussed below.

¹ Dept. of Physics & Astronomy
University College London
Gower Street
London WC1E 6BT

2 WR photometric systems

The standard Johnson UBV filter set provides adequate continuum measurements for Luminous Blue Variables and slash stars. However, the intrinsic colors of W-R stars are complicated by their strong emission lines. These features greatly modify the flux measured in each optical filter, especially those of the relatively broad band UBV system. Such measurements may overestimate the true continuum level by up to 1 magnitude in extreme cases, or more typically 0.5 mag for single early-type W-R stars. Consequently, Smith (1968b) introduced *narrow band ubv* filters specifically designed for W-R stars to minimize the effect of emission lines (but their effect cannot be entirely eliminated). Most photometry of W-R stars has used this *ubv* filter system.

The following relations relate the broad and narrow band optical indices for W-R stars (from van der Hucht 2001).

$$\begin{aligned} E_{B-V} &= 1.21 E_{b-v} \\ A_v &= 4.1 E_{b-v} = 1.11 A_V \\ (B - V)_0 &= 1.28 (b - v)_0 = 0.11 \\ M_V &= M_v + 0.1 \end{aligned}$$

A recent compilation of the spectral subtype dependence of $(b - v)_0$ for W-R stars (van der Hucht 2001) reveals values in the range -0.2 to -0.35 for WN and WC stars, with the possible exception of WC9 stars. Observations in the *ubv* system are mostly used to determine the interstellar extinction, A_v , rather than to be able to say anything about the star itself.

As with OB stars, absolute magnitudes of slash stars and W-R stars rely on calibrations of stars that are members of Galactic or Magellanic Cloud clusters or OB associations. Confirmed numbers of LBVs are insufficient for absolute magnitudes to be useful. Only γ Velorum (WC8+O), the only naked eye W-R star visible (from the southern sky) was close enough for a reliable Hipparcos measurement.

van der Hucht (2001) also provides a modern M_v -spectral type calibration for WN and WC stars. Absolute visual magnitudes for single W-R stars range from $M_v = -2.5$ for early WN and WO stars, to $M_v = -4.5$ for late-type WC stars, and $M_v = -7$ for late-type WN stars. There are no reliable bolometric corrections published for W-R stars to date, although individual values range from -2.7 amongst very late WN stars (Crowther & Smith 1997), to approximately -6 for weak-lined, early-type WN stars (Crowther et al. 1995c) and WO stars (Crowther et al. 2000). For LBVs and slash stars, bolometric corrections closely mimic those of corresponding OB supergiants (Crowther 1997).

3 Spectral Classification

3.1 W-R stars

Spectral classification of W-R stars is derived from emission line strengths and line ratios (Beals & Plaskett 1935). The modern system for both nitrogen-rich (WN)

and carbon-rich (WC) stars is based upon that introduced by Smith (1968a). Details of the classification scheme are contained in van der Hucht (2001) where an up-to-date catalogue of Galactic W-R stars is also given.

WN spectral subtypes utilize a one-dimensional scheme, involving line ratios of N III-V and He I-II lines, ranging from WN2 to WN9, extended to later subclasses by Smith et al. (1994) to accommodate cool slash stars (see below). At the earliest classical subtypes solely He II and N V are present, with He I and lower ionization stages of nitrogen absent, whilst at the latest subtypes, He I is strong, He II present but weak, N III present, with higher ionization stages absent. Since the definition of some subtypes (e.g. WN8) involves the relative strength of nitrogen to helium emission lines, there may be an inherent metallicity dependence. Consequently, Smith et al. (1996) devised a scheme that involved ratios of lines from either helium or nitrogen. Various multi-dimensional systems have been proposed, generally involving line strength or width, none of which have been generally adopted, the most recent of which was by Smith et al. (1996), including criteria for the presence of hydrogen (h), and broad (b) line widths.

WC spectral subtypes depend on the line ratios of C III and C IV lines along with the appearance of O V (Smith et al. 1990), spanning WC4 to WC9 amongst massive stars ¹, which has recently been revised to account solely for C III-IV lines (Crowther et al. 1998). Oxygen-rich WO stars form an extension of the WC sequence to higher excitation, exhibiting strong O VI emission (Barlow & Hummer 1982). The most recent scheme involves WO1 to WO4 depending on the relative strength of O V-VI and C IV emission lines (Crowther et al. 1998). Although line widths of early WC and WO stars are higher than late WC stars, width alone is not a defining criterion for each spectral type.

As with O stars, atlases of W-R spectra are available at ultraviolet (Willis et al. 1986) and near-IR (e.g. Figer et al. 1997) wavelengths, permitting approximate spectral types to be assigned.

3.2 *Slash Stars*

As described in my first article, a subset of O stars exhibit emission lines in their optical spectra. Although these are generally much weaker than in W-R stars, there are a number of stars with intermediate characteristics, which have been coined ‘transition’ objects between O-type and W-R. There are two flavours, hot and cool ‘slash’ stars.

3.2.1 O3If/WN Stars

Walborn (1973) discussed the close morphological relationship between various early-type stars in Carina, namely HD 93129A (O3 If, recently revised to O2 If

¹Some H-deficient central stars of Planetary Nebulae show carbon-sequence W-R spectral features, that extend to later spectral types (Crowther et al. 1998).

by Walborn et al. 2002) and HD 93162 (WN6ha). Subsequently, stars with intermediate characteristics were identified in the LMC, such as Sk-67° 22 and Melnick 42, and assigned spectral types of O3If/WN (Walborn 1986). Whatever the nomenclature issues, the important point is that merely a small change in wind density distinguishes the most extreme O2–3 supergiants from the least extreme WN stars, such as those identified in Carina, R136 and NGC 3603. These WN stars are hydrogen-rich (h) and contains (apparently) intrinsic absorption (a) lines (Crowther et al. 1995b; Crowther & Dessart 1998), such that they have apparently more in common with early O supergiants than the bulk of hydrogen-free, core He-burning W-R stars.

For completeness, cooler analogues of the hydrogen-rich Carina WN stars have also been identified. A careful reconsideration of the spectral types of three Galactic O6–8Iafpe stars led Bohannan & Crowther (1999) to revise their classifications to WN9ha. They provided suitable criteria for consideration of previously undiscovered stars with similar spectral morphologies.

3.2.2 Ofpe/WN9 Stars

Walborn (1977, 1982) identified a group of late O LMC emission line stars as Ofpe/WN9 subtypes, with apparently intermediate characteristics of Of and late WN stars. Members of this group were re-classified WN9, 10 or 11 by Smith et al. (1994) and Crowther & Smith (1997) on the basis that characteristic O star photospheric lines were absent. Additional examples have been identified in M33, IC 10 and NGC 300. WN11 subtypes closely resemble extreme early-type B supergiants, except for the presence of He II $\lambda 4686$ emission. Interest in this group of very late WN stars grew because of their high frequency of associated H II regions, plus an apparent connection with LBVs. One prototypical Ofpe/WN9 star (subsequently re-classified as WN11), R127 in the LMC, was later identified as a LBV. Conversely, AG Car, a well known Galactic LBV, exhibited a WN11-type spectrum at extreme visual minimum (Smith et al. 1994).

In the K -band, WN9–11 stars exhibit strong He I $2.058\mu\text{m}$ and Br γ emission lines (Figer et al. 1997; Bohannan & Crowther 1999). Observations obtained within the last decade reveal that a large number of stars within clusters at, or close to, the Galactic Centre also show this spectral morphology. As in the optical, very late WN stars also resemble early B supergiants, such as P Cygni. Consequently, the presence of He II near-IR lines is required to confirm these stars as bona-fide WN stars, which is currently lacking in most cases. Examples are shown by Najarro et al. (1997a) for He I sources in the central cluster, Figer et al. (1999a) in the Quintuplet cluster Emission-line stars located in the Arches cluster most closely resemble hot slash stars or weak-lined WN stars (Blum et al. 2001).

3.3 Luminous Blue Variables

Observationally, LBVs inhabit a part of the H-R diagram that is bereft of normal stars. This region is known as the Humphreys-Davison limit (Humphreys &

Davidson 1979), above which either no supergiants are observed, or their transit through this region is so brief that none would be expected on statistical grounds. The photometric and spectroscopic variability of LBVs is suspected of being so large since they stars lie close to the Eddington limit. Beyond this limit, stars would not remain stable.

In contrast with O and W-R stars, there is no single spectroscopic defining characteristic to define a LBV; historically it simply had to be a variable, luminous, blue star (Conti 1984). Examples are known in our Galaxy, including η Carinae and P Cygni, the Magellanic Clouds, known collectively as S Doradus-type variables, plus M31 and M33, these known historically as the Hubble-Sandage variables, after their discoverers in the 1950's. More recently, it has been proposed that the presence of an associated H II region, a remnant from recent mass-losing episodes, may be a defining characteristic.

LBVs are generally restricted to those stars with substantial photometric variability on timescales of decades, plus associated spectroscopic changes. Two types of photometric variability are commonly discussed.

3.3.1 LBVs during quiescence

LBVs exhibit micro-variations (~ 0.1 mag) over short timescales, and irregular variations of up to 1–2 mag, in which the spectral type may vary between an early B supergiant at visual *minimum* to a late B or early A supergiant at visual *maximum*. Indeed, some LBVs have even proceeded to spectral types as late as F or even G. Classic examples of irregular variability over the past few decades include AG Car and HR Car in the Milky Way, R 71 and R 127 in the LMC.

Because of the irregular nature of variability, it is almost certain that many ‘dormant’ LBVs await discovery, with all luminous blue supergiants potentially undergoing a LBV outburst at some point in their evolution. A fine example is HDE 316285, an early B supergiant with a more extreme wind than P Cyg and spectroscopic similarities to η Car (Hillier et al. 1998). Only moderate spectroscopic variability has been observed to date, such that an LBV nomenclature is not (yet) appropriate.

3.3.2 Giant eruptions

More remarkably, giant eruptions in LBVs have been observed, in which photometric variability exceeds 2 mag in the visual. Such outbursts are exceedingly rare, with only P Cygni and η Carinae known to have undergone such an eruption between the 15th and late 20th centuries. It is almost certain that other stars will have undergone such eruptions during this time, but were not sufficiently bright to have been noticed by astronomers. P Cygni was discovered in 1600 by the Dutchman Bleau, when it suddenly appeared as a naked eye star (no star was apparent before). It remained bright for many years, before fading and re-appearing in 1655, and then fading again. At present it is around 5th magnitude, and so is at present the brightest LBV in the sky. P Cygni profiles, generally observed in UV

resonance lines in hot star winds are described as such because P Cygni displays these profiles in hydrogen and helium lines in its optical spectrum.

During the early 1990's, two other stars have erupted as LBVs – the well known WN+O binary (or perhaps triple) system HD 5980 in the Small Magellanic Cloud (Barba et al. 1995), plus a previously unknown star in the GH II region NGC 2363 of the low-metallicity galaxy NGC 2366, labelled V1 (Drissen et al. 2001).

3.4 B[e] supergiants

The term ‘B[e] star’ was coined by Conti (1976), and designates those B stars which show forbidden (‘[]’) emission (‘e’) lines in their optical spectrum. B[e] stars should not be confused with main-sequence Be stars. B[e] supergiants, or sgB[e] following Lamers et al. (1998), are extremely luminous massive stars observed in the Magellanic Clouds that are distinguished by a plethora of low-excitation emission lines including (broad) Balmer lines and (narrow) singly ionized metals, including permitted and forbidden iron lines. These stars also show a significant near- and mid-IR dust excess, providing evidence for hot, circumstellar dust, such that the standard interpretation for these unusual properties is an equatorial excretion disk, plus a fast polar stellar wind (Zickgraf et al. 1985). In common with other luminous supergiants, B[e] supergiants appear to be nitrogen rich, showing evidence for the products of the CN-cycle at their surfaces. Gummertsbach et al. (1995) provide a thorough discussion of the luminosities and effective temperatures of the known B[e] stars in the Magellanic Clouds.

B[e] supergiants occupy a similar location in the H-R diagram to the LBVs, although a firm connection is currently lacking (Conti 1997). It is likely that such stars are not post-red supergiants, since angular momentum would be lost during this phase which would not be regained once the stars return blueward. Photometric variability of B[e] supergiants is generally small, although the binary system R4 in the SMC has shown an LBV-type photometric and spectroscopic outburst of 0.5 mag in the mid-1980s (Zickgraf et al. 1996). Hen S18, another SMC B[e] supergiant is also spectroscopically variable (Sanduleak 1977),

4 Circumstellar nebulae

The recent mass-loss history of a subset of hot, luminous stars may be studied their circumstellar environment.

η Car, at the heart of the great Carina nebula, is known to have shed approximately $3M_{\odot}$ during a couple of decades between 1837–1860. During that time it became one of the brightest stars in the southern sky and so has one of the most remarkable photometric histories of any naked-eye star. The event remains unprecedented to this day, and corresponds to losing the equivalent mass of the Earth every 15 minutes for 2 decades. Such a large amount of material should be visible. Indeed it is, and has been named the Homunculus nebula. HST/WFPC2 images of the Homunculus reveal incredible bi-polar emission lobes (Morse et al. 1998). Its orientation on the sky is such that the central star(s) is largely masked

from view. Damineli et al. (1997, 2000) found evidence in favour of a 5.5 year period for η Car involving two massive $\sim 70M_{\odot}$ stars.

Nevertheless, we do see evidence that other LBVs underwent similar eruptions through their circumstellar environment. The ejecta nebula associated with the famous Luminous Blue Variable AG Car probably differs only from the Homunculus Nebula through a much greater dynamical age and orientation (Langer et al. 1999). The dynamical age of a nebula, τ_{dyn} , is simply the radius of the nebula, divided by the expansion velocity. The size of the nebula associated with AG Car is 0.6 pc (assuming a distance of 6 kpc), whilst the expansion velocity is 70 km s^{-1} such that $\tau_{\text{dyn}} = 8,400 \text{ yr}$, and is typical of other LBVs, except for η Car, for which its dynamical age is in good agreement with the 1837 event.

The Pistol Star at the Galactic Centre represents the best example of an early-type star that has not been observed to dramatically vary in continuum flux, but possesses a H II region (Figer et al. 1998, 1999b), a characteristic which has been used to classify it as an LBV.

Ejecta H II regions are also associated with a subset of Wolf-Rayet stars, predominantly amongst WN stars. Beautiful examples include M1 67 (Grosdidier et al. 1998) and NGC 6888 (Moore et al. 2000). It is likely that these were ejected during a previous (LBV?) phase, but are currently photo-ionized by the W-R star. Other W-R stars possess associated H II regions, albeit composed of material exhibiting ISM abundances (Esteban et al. 1992).

5 Stellar Properties

As discussed in my first article relating to OB stars, spectral line techniques, as opposed to continuum techniques, are required to derive stellar properties of W-R and other early-type mass-losing stars. Such rules apply equally to LBVs during their A and B-type spectral phases, since they too possess strong stellar winds, and their surfaces are probably H-depleted.

Spherically extended, non-LTE line blanketed model atmospheres are essential for the detailed analysis of Wolf-Rayet stars, slash stars and Luminous Blue Variables, of which CMFGEN (Hillier & Miller 1998) and that developed by the Potsdam group (Gräfener et al. 2002) have been most widely used to date.

5.1 Temperatures

As with normal O stars, spectral lines of helium (or nitrogen) may be used to derive the T_{eff} of Ofpe/WN9 stars and WN stars, whilst silicon diagnostics offer the possibility of determining temperatures of LBVs during their B supergiant phases. Carbon or oxygen lines can be used to diagnose WC and WO stars. Stellar temperatures for spherically extended stars are more difficult to characterize than dwarfs, since the geometric extension in these stars is comparable with the stellar radius particularly for W-R stars and LBVs. In practice, one defines a radius as the distance from the center where the mean Rosseland optical depth, τ_{Ross} , is of order 10. For normal stars this does not differ from the usual radius at $\tau_{\text{Ross}} = 2/3$, but

differs greatly for W-R stars, and represents a much more representative ionization indicator. In extreme cases, such as η Car, the high density of the wind completely veils the stellar photosphere such that it is impossible to characterize a stellar temperature (Hillier et al. 2001).

We have already discussed that the recent incorporation of line blanketing in analysing the *photospheric* spectrum of OB stars has led to a substantial *decrease* in derived temperatures (Martins et al. 2002; Crowther et al. 2002b). In contrast, similar advances for W-R stars has led to an *increase* in derived temperatures obtained from *emission* lines (Herald et al. 2001). Stellar temperatures of W-R stars range from 30,000K amongst late-type WN stars to well in excess of 100,000K for early-type WN stars and WO stars, although differences in wind density complicate accurate determinations. Hot slash stars and H-rich WN stars have stellar temperatures at the high end of the O main-sequence, typically $\sim 40\text{--}45\text{kK}$. These stars have the highest luminosities of all ‘normal’ stars.

Increased stellar temperatures of W-R stars relative to OB stars implies that their contribution to the Lyman continuum of young stellar clusters is increased. However, since blanketing redistributes extreme-UV radiation to lower energies, predicted fluxes at higher energies, such as those shortward of the He II $\lambda 228$ edge, are actually reduced (Smith et al. 2002). It is solely hot W-R stars with relatively weak winds that are a significant source of He II Lyman continuum radiation.

Luminous Blue Variables range in stellar temperatures between $\sim 8\text{kK}$ at visual maximum, when they typically exhibit an A-type hypergiant spectrum, and 25kK at visual minimum, with an early B subtype. Consequently these stars are minor contributors to the Lyman continuum photons of young clusters. Because of the change in B.C. with T_{eff} , visual changes in LBVs are generally considered to occur at fairly *constant* bolometric luminosity (Crowther 1997). It is only for LBVs undergoing giant eruptions, such as η Car in the 19th Century, that this is not true. Finally, WN9-11 stars, possess temperatures in the range $25\text{--}30\text{kK}$, filling-in the temperature sequence between LBVs and W-R stars (Smith et al. 1994; Crowther & Smith 1997).

5.2 Masses

Spectroscopic measurement of masses in post-main sequence, early-type stars is possible in principle via analysis of line wings. However, stellar winds in most cases affect line spectra sufficiently to hinder gravity determinations. Similarly, evolutionary masses are often complicated by the uncertain (blueward or redward) evolutionary state of the star. Estimates of initial masses can be obtained from cluster turn-offs (e.g. Massey et al. 2001; St-Louis et al. 1998).

In the case of Wolf-Rayet stars and Luminous Blue Variables, one relies on direct measurements of current masses via analysis of binary orbits. Amongst LBVs, only R81 in the LMC is a binary with a well characterized orbit (Tubbesing et al. 2002), whilst several examples are known amongst W-R stars, widely studied by V. Niemela and colleagues. By way of example, the binary nature of γ Velorum permits a mass determination for both the W-R and O components, with

$10M_{\odot}$ obtained for the WC component. Fortunately, there is predicted to be a robust relationship between the mass of a hydrogen-free WN or WC star and its stellar luminosity (Schaerer & Maeder 1992). For γ Vel, good agreement is found between the direct measurement and that resulting from a recent line-blanketed spectroscopic analysis (De Marco et al 2000).

Not all W-R stars have masses which are relatively low. The eclipsing binary Carina WN7ha+O system HD 92740 has provided an exceptionally high mass for the WN component of $\sim 55M_{\odot}$ (Schweickhardt et al. 1999). This result, together with spectral analyses revealing that this star is rich in hydrogen has led to a subset of W-R stars being dubbed H-burning main-sequence O stars with particularly strong stellar winds. Unfortunately, direct information on the processes going on within the core of a massive star are very hard to determine without recourse to evolutionary models.

6 Stellar Wind Properties

As for OB stars, several techniques may be used to study the wind properties of W-R stars and LBVs. These stars have the advantage that their more powerful winds permit velocity and mass-loss rate determinations other than in the UV and at $H\alpha$. Analytical relationships (Wright & Barlow 1975) permit measurements of mass-loss rates of LBVs and W-R stars via IR-radio continua, providing distances and wind velocities are known. Current sensitivities of radio telescopes restricts this method to stars within a few kpc of the Sun.

6.1 Wolf-Rayet stars

In addition to the usual ultraviolet P Cygni wind velocity diagnostics (Prinja et al. 1990), optical, and especially near-IR P Cygni lines of He I permit accurate wind velocities for W-R stars (Howarth & Schmutz 1992; Eenens & Williams 1994). ISO mid-IR fine structure lines also facilitate direct measurement of wind velocities in some cases. Wind velocities of WR stars increase from late, to early-spectral types. WN8 and WC9 stars possess wind velocities of order $\sim 1000 \text{ km s}^{-1}$, whilst WN3 and WC4 stars have winds on average a factor of three times larger. WO stars possess the highest wind velocities of any (stable) star, up to 5500 km s^{-1} (Kingsburgh et al. 1995).

Radio mass-loss rates of W-R stars have been derived by Leitherer et al. (1997) using the Wright & Barlow (1975) analytical relationship, although care needs to be taken with regard to the elemental abundances and ionization balance in the radio emitting region. This method is distance limited, due to the limited sensitivity of current radio telescopes, and one should obtain radio fluxes at more than one wavelength in order to avoid overestimates of mass-loss rates due to non-thermal emission in a close binary (Chapman et al. 1999). Non-LTE model analysis of UV, optical and IR line profiles is more widely used for Galactic and extragalactic W-R stars (e.g. Dessart et al. 2000).




Fig. 1. Comparison between (clumped) mass-loss rates of Galactic (open symbols) and LMC (filled symbols) WC stars, revealing systematic weaker winds in the latter, by an amount predicted by radiatively driven winds. The solid line is a fit to the LMC data (Crowther et al. 2002a), whilst the dotted line is a calibration from Nugis & Lamers (2000).

Unblended spectral regions are scarce in the ultraviolet and visual, particularly for WC stars, such that synthesis of fluxed spectroscopy is the preferred method. It quickly became apparent that it was impossible to reproduce both peak line intensities – the strength of which are proportional to the square of the density – and red electron scattering winds – linearly proportional to the density – under the assumption of homogeneity (Hillier 1991). Many other observations (Lepine et al. 2000) and theoretical arguments (Dessart & Owocki 2002) point to severely clumped winds in W-R stars, although precise volume clumping factors are challenging to derive.

Most observations are consistent with clumping factors of $f \approx 0.1$, in the sense that clumped mass-loss rates are a factor of $1/\sqrt{f} \approx 3$ times lower than previous homogeneous rates (Hamann & Koesterke 1998). Clumped W-R mass-loss rates typically span a range from $1\text{--}3 \times 10^{-5} M_{\odot} \text{ yr}^{-1}$, such that wind performance numbers, $\dot{M}v_{\infty}/(L/c)$ the factor by which the wind exceeds the single scattering limit, lie in the range 1–10. Nugis & Lamers (2000) provide calibrations for WR mass-loss rates as a function of luminosity, He and metal content.

6.2 *Metallicity dependence for WR winds?*

In contrast with normal OB stars, it is not yet established that W-R winds are (solely) driven by radiation pressure. A metallicity dependence of O star winds is well established theoretically (e.g. Vink et al. 2001), for which observational support has been demonstrated (Puls et al. 1996).

In the case of W-R stars, derived mass-loss rates of Milky Way WN stars span a wide range, such that comparisons with Magellanic Cloud WN stars can be interpreted in a variety of ways (Crowther & Smith 1997; Crowther 2000). In contrast, the wind densities of WC stars form a much cleaner distribution. Crowther et al. (2002a) demonstrated that single early-type WC stars in the LMC possessed winds that were weaker than those of comparable Milky Way stars. This difference, identical to that expected from radiatively driven wind theory for OB stars, led Crowther et al. (2002a) to conclude that WC spectral types result primarily from metallicity differences, rather than abundances as suggested by Smith & Maeder (1991).

Rotation is much more difficult to measure in W-R stars than in OB stars, since photospheric features are lacking. Attempts have been made, but without much success. Spectropolarimetry does permit information to be obtained regarding geometry, which indicates that most Galactic W-R stars are spherical, with a few notable exceptions amongst WN stars (Harries et al. 1998).

6.3 *Slash stars*

Hot slash stars (O3If/WN stars) have wind properties intermediate between O2–3 supergiants, and weak-lined, hydrogen rich WN5–7 stars, i.e. high wind velocities of $\sim 3000 \text{ km s}^{-1}$, plus mass-loss rates approaching $10^{-5} M_{\odot} \text{ yr}^{-1}$ (de Koter et al. 1994). In contrast, cool slash stars (WN9–11 stars) possess relatively low wind velocities, generally $< 500 \text{ km s}^{-1}$, plus moderate mass-loss rates (Crowther et al. 1995a; Crowther & Smith 1997). Galactic Centre slash stars have higher wind velocities, in the range $500\text{--}1000 \text{ km s}^{-1}$ (Najarro et al. 1997a).

6.4 *Luminous Blue Variables*

Typically, LBVs during quiescence possess wind velocities which are a factor of several times lower than corresponding OB supergiants, with mass-loss rates a factor of several times higher (Crowther 1997). For example, P Cygni has a wind velocity of 185 km s^{-1} , in contrast to a typical $v_{\infty} = 1060 \text{ km s}^{-1}$ for normal B1 supergiants (Prinja et al 1990). Leitherer (1997) studied the mass-loss variability of AG Car during its excursions around the H-R diagram, deducing a ‘random walk’ in which the mass-loss rate did not undergo dramatic variations. R71 appears to be unusual in possessing a much reduced mass-loss rate at visual minimum (Leitherer et al. 1989).

In contrast, LBVs undergoing a major eruption possess exceptional wind properties. η Car is believed to have shed in excess of $3M_{\odot}$ over between 1840–1860, such that its average mass-loss rate during that event was $\sim 10^{-1} M_{\odot} \text{ yr}^{-1}$. Even

today, during its quiescent phase, its mass loss rate is perhaps only a hundred times lower. Note that a giant eruption does not necessarily imply exceptional mass-loss. NGC 2363-V1 is an LBV that is currently undergoing a giant eruption with a mass-loss rate that is comparable to η Car during *quiescence* rather than *eruption* (Drissen et al. 2001).

7 Abundances

7.1 Wolf-Rayet stars

It has long been suggested that W-R abundances represented the products of core nucleosynthesis. However, it was only with Balmer-Pickering decrement studies (Conti et al. 1983) that indeed hydrogen was found to be severely depleted in all WR stars. Nitrogen and carbon abundances are consistent with CNO cycle values in WN stars from both atmospheric and nebular analyses, with nitrogen representing up to $\sim 1\%$ of the surface mass. In WC and WO stars extremely high carbon and oxygen abundances are found, typically in excess of 50% by mass, clearly reflecting both triple- α and α -capture products.

Recombination line studies (Smith & Hummer 1988; Smith & Maeder 1991), using theoretical coefficients for different transitions are most readily applicable to WC stars, since they show a large number of spectral lines in their optical spectra. Atomic data is most readily available for hydrogenic ions, such as C IV and O VI, so early-type WC and WO stars can be most easily studied. In reality, optical depth effects play a role too. Limitations with earlier approaches have greatly been overcome with modern spectroscopic non-LTE studies. WN abundance patterns are consistent with material processed by the CNO nuclear burning cycle, whilst WC stars reveal products of He burning.

A clear subtype effect in the H-abundance is found for WN stars, with late-type WN stars generally showing some hydrogen, with early-type WN stars hydrogen free. Exceptions to this general rule do exist, such as HD 177230 (WN8o, Crowther et al. 1995b) and HD 187282 (WN4(h), Crowther et al. 1995c). WNLha stars are rich in hydrogen (Crowther & Dessart 1998). Non-LTE model atmosphere analyses indicate fully CNO processed material at the surfaces of most WN stars (e.g. Crowther et al. 1995b).

Conti & Massey (1989) identified 7 Galactic and 2 LMC stars as having intermediate WN–WC characteristics based on anomalously high C IV $\lambda 5801$ /He II $\lambda 4686$ line strengths compared to normal WN stars. Modern analyses of these ‘transition’ WN/C stars indicate elemental abundances intermediate between WN and WC stars, with N/C ~ 1 by number (Crowther et al. 1995d), supporting rotationally induced mixing in W-R stars.

Neither hydrogen nor nitrogen is detected in the optical, infrared or ultraviolet spectra of WC stars. Early recombination line studies of WC stars indicated a trend of increasing C/He from late- to early- WC subtypes (Smith & Hummer 1988) with C/He = 0.04–0.7 by number, clearly indicative of the exposition of He-burning products. More recent, non-LTE model studies indicate C/He = 0.1–0.5

by number, with no such subtype trend (Koesterke & Hamann 1995). Indeed, LMC WC stars possess similar surface abundances to Milky Way counterparts Crowther et al. (2002a). Oxygen abundances are more difficult to determine, although values rarely exceed $O/H=0.1$ (Dessart et al. 2000; Crowther et al. 2002a). Neon is predicted to be enhanced in WC atmospheres, as supported by results based on mid-IR fine-structure lines from ISO (Dessart et al. 2000), once the clumped nature of W-R winds are taken into account. WO stars are extremely carbon and oxygen rich, as deduced by recombination analyses (Kingsburgh et al. 1995) and non-LTE models (Crowther et al. 2000), with $C/He \geq 0.5$ and $O/He \geq 0.1$ by number.

7.2 *Slash stars*

All WN9–11 stars discovered to date still possess hydrogen at their surface, albeit substantially reduced, with $H/He=1.2$ – 3.5 by number for those optically visible, and $H/He \leq 2$ for Galactic Centre He I sources (Najarro et al. 1997a). These also exhibit partially processed CNO material from both stellar (Crowther et al. 1995a; Crowther & Smith 1997) and nebular (Smith et al. 1998) studies. Abundances patterns of O3If/WN stars mimic those of hydrogen rich WNha stars, in the sense that H is little depleted, if at all, whilst CNO elements are partially processed (Heap et al. 1991; de Koter et al. 1997).

7.3 *Luminous Blue Variables*

Crowther (1997) compares surface He-contents in five LBVs, revealing a surprisingly uniform mass fraction of $H/He=2$ – 2.5 by number. Najarro et al (1997b) has obtained $H/He \sim 3$ from a detail investigation into P Cygni. CNO abundances in LBVs have been rarely studied directly (e.g Lennon et al. 1997). More frequently, these are studied via analysis of the N and O abundances from their associated H II regions (Smith et al. 1998). Lamers et al. (2001) discuss evolutionary implications for CNO abundances determined in LBV nebulae.

Most recently, high quality UV and optical spectra of η Car have been obtained with HST and analysed to reveal enriched nitrogen by at least a factor of ten, whilst carbon and oxygen are depleted (Hillier et al. 2001).

8 W-R stars in Local Group galaxies

The presence of strong emission lines in W-R stars permits their discovery at large distances via use of suitable narrow-band filters. For example, a set of narrowband filters centred at $\lambda 4650$, $\lambda 4686$ and a nearby continuum region will discriminate between WC, WN and normal stars. Groups led by P. Massey, and by A.F.J. Moffat in the 1980's used such filter combinations to find suitable W-R candidates in the Milky Way and other Local Group galaxies. Suitable filter sets on large 8–10m telescopes now facilitates identification of W-R stars beyond the Local Group (e.g. Schild et al. 2003).




Fig. 2. WC/WN ratio for Local Group and Sculptor Group spiral (squares) and irregular galaxies (circles) versus oxygen content. Individual data points are taken from Massey & Johnson (1998), Schild et al. (2003), except for IC 10 for which recent spectroscopic (Crowther et al. 2003, filled circle) and photometric (Massey & Holmes 2002, open circle) results are indicated.

8.1 Visibly obscured W-R stars within the Milky Way

Since massive stars are solely formed within the disk of the Milky Way, only a small number can be observed at optical or ultraviolet wavelengths. At present, little more than 200 W-R stars are known within our Galaxy, yet many thousand are expected to be present (van der Hucht 2001). It is only via observation at IR wavelengths where interstellar extinction is much reduced that one can detect individual stars. We have already discussed the young massive clusters close to the centre of our Galaxy, which contain hot and cool slash stars, plus several normal W-R stars. Other Galactic clusters, of which Westerlund 1 is a beautiful example, have recently been found to host large W-R populations. Clark & Negueruela (2002) have identified at least 11 new W-R stars in Westerlund 1, plus a wide diversity of blue, yellow and red supergiants, such that it has a mass comparable with any young cluster in the Local Group.

Near-IR surveys, using filters selected to measure W-R emission lines and continuum regions, are presently underway (e.g. Homeier et al. 2003). Real progress awaits application of this method using large format detectors mounted on large 4m-class telescopes.

8.2 WR subtype distribution – the puzzling case of IC 10

It is well known that the distribution of W-R stars, specifically the ratio of W-R to O stars, is a strong function of environment. One can understand this general behaviour from the metallicity dependence of wind strengths prior to the W-R phase, in the sense that weaker winds at low metallicities, such as in the SMC, cause lower mass-loss during the main and immediate post-main sequence, such that higher initial mass is needed for the formation of a W-R star via single star evolution (Meynet & Maeder 1994).

In addition to the reduced W-R/O fraction at lower metallicity, it is also well known that the number of WC stars to WN stars is also a sensitive function of metallicity in Local Group galaxies. In the Milky Way, WC/WN=3/4 (van der Hucht 2001), versus WC/WN=1/10 in the SMC (Massey & Duffy 2001). Fig. 2 compares WC/WN ratios with metallicity for various Local Group and Sculptor Group galaxies.

Late WC subtypes are restricted to metal-rich environments, such as the inner Milky Way and M31, with early WC stars found predominantly in metal-poor regions, such as the Magellanic Clouds. Crowther et al. (2002a) attributed this difference to weaker WC winds in metal poor regions, since the C III $\lambda 5696$ classification line is very sensitive to wind density.

Massey (1996) first highlighted the apparent anomaly of IC 10, which is unique amongst Local Group galaxies, in the sense that it is metal weak, but has a large number of WR stars, plus an uniquely high WC/WN ratio of ~ 2 (Massey & Johnson 1998). Recent spectroscopy of candidates that had been identified by Massey et al. (1992) and Royer et al. (2001) confirmed a high WC/WN ratio of 1.3 (Crowther et al. 2003). All WC stars were early subtypes, with one exception, contradicting photometric claims of several late WC stars in IC 10 (Royer et al. 2001).

Recently, a new imaging survey has been carried out by Massey & Holmes (2002) suggesting that IC 10 possesses a normal WC/WN ratio for its metallicity, such that many dozens of (principally) WN stars had not been detected by earlier surveys. Of course, spectroscopic follow up is needed to verify the nature of new candidates. Whatever the case, it appears that IC 10 has an exceptional massive star content and represents the closest galaxy-wide starburst.

W-R stars have been observed in very metal deficient star forming galaxies, such as I Zw 18 (de Mello et al. 1998), and also also seen in in star forming galaxies at high redshift (Shapley et al. 2003).

9 Cool Supergiants

Up to now, this article has dealt with early-type massive stars. Many, if not all, massive stars briefly pass through a yellow (FG) or red (KM) supergiant phase, of which Betelgeuse, α Ori (M2.2 Iab) is the best known example. Moderately massive stars end their life in the red as a Type II supernova (see later), whilst very massive stars either circumvent the cool supergiant phase entirely, or subsequently

return to the blue as a W-R star. Surprisingly few clusters are known in which both red supergiants (RSG) and W-R stars are known to co-exist. Collinder 228 and Berekley 87 each contain one RSG and W-R star (Massey et al. 2001). Remarkably, Westerlund 1 hosts at least 2 RSG and 11 W-R stars (Clark & Negueruela 2002).

Mass-loss in cool supergiants is episodic and involves slow, dense outflows. Radial pulsations are generally assumed to provide the mechanism of initiating mass-loss in M supergiants, as evidenced from circumstellar dust shells (Bowers et al 1983), with some notable exceptions – VY CMa (M5 Ia) and ρ Cas (F8 Ia+). Outflow velocities of RSG are typically $10\text{--}40\text{ km s}^{-1}$ (Jura & Kleinmann 1990).

Since we are concerned with massive stars here, we shall restrict our discussion to the most luminous ‘hypergiants’ (de Jager 1998) with $\log(L/L_{\odot}) > 5$. Studies have largely focussed on M supergiants, with only highly unusual FG supergiants studied in detail. IRC+10420 in particular is widely interpreted as a RSG rapidly moving blueward (Jones et al. 1993; Humphreys et al. 2002), which has recently experienced huge variability in mass-loss, 10^{-2} to $10^{-4}\text{ }M_{\odot}\text{ yr}^{-1}$, via analysis of its circumstellar dust shell (Blöcker et al. 1999).

Various techniques have been used to quantify mass-loss in late-type supergiants, whose wind driving mechanism remains uncertain, although these remain much more controversial than for early-type stars. For example, recent mass-loss estimates for VY CMa span a factor of twenty – $1.6 \times 10^{-5}\text{ }M_{\odot}\text{ yr}^{-1}$ (Josselin et al. 2000) to $3.5 \times 10^{-4}\text{ }M_{\odot}\text{ yr}^{-1}$ (Stanek et al. 1995).

Most widely, sub-mm CO emission or mid-IR dust emission permit mass-loss estimates for cool supergiants. Jura & Kleinmann (1990) and Reid et al. (1990) have estimated mass-loss rates of Galactic and LMC RSG, respectively, in which the LMC sample formed a tight group with $\sim 10^{-4.5}\text{ }M_{\odot}\text{ yr}^{-1}$ and the Galactic sample spanned a wide range from 10^{-6} to $10^{-4}\text{ }M_{\odot}\text{ yr}^{-1}$. More recently, Sylvester et al. (1998) obtained *higher* mass-loss rates of 10^{-5} to $10^{-4}\text{ }M_{\odot}\text{ yr}^{-1}$ for Galactic M supergiants using the $10\mu\text{m}$ silicate emission feature, whilst van Loon et al. (1999) have obtained LMC RSG mass-loss rates of $\leq 10^{-5}\text{ }M_{\odot}\text{ yr}^{-1}$, *lower* than Reid et al. Therefore, depending on the studies used, LMC supergiants either exceed those in the Milky Way, or vice versa. It is clear that much research remains to be undertaken in order to firmly establish mass-loss in late-type supergiants.

10 End states of massive stars

10.1 Supernovae

All supernovae (SN), with the exception of Type Ia’s, originate from the core collapse of massive stars. Those ending their life as red supergiants explode as a Type II SN, although in some rare cases, Type II SN progenitors are blue supergiants, such as Sk-69° 202, the B3I progenitor of SN1987A. The origin of this remains unclear, although the most credible scenarios involve binarity. Stars which were initially even more massive end their life as either a Type Ib or Ic SN. These are presumably related to WN and WC progenitors, respectively, given that neither

possess spectral lines of hydrogen, whilst helium is also absent in Type Ic's. Considering all galaxy types in the Local Universe, Cappellaro et al. (1999) found that 70% of all observed SN are due to core collapse massive stars, predominantly Type II.

Light curve and spectroscopic analysis of SN permit ejecta masses to be determined. Iwamoto et al. (1998, 2000) derived ejecta CO masses of $2\text{--}14M_{\odot}$ for the Type Ic supernovae SN 1994I, SN 1997ef and SN 1998bw. The masses discussed here represent minimum masses for the final progenitor, since an unknown (model dependent) fraction would have imploded to form a black hole or other compact object. Nevertheless, the ejecta masses of SN 1997ef and SN 1998bw fit in well with the final masses of LMC WC stars derived by Crowther et al. (2002a).

SN 1998bw possessed exceptional wind velocities ($27\,000\text{ km s}^{-1}$) immediately after the explosion, such the energy of the ejecta exceeded 10^{52} erg. Consequently, it has been denoted a 'hypernova', an exceptional class of energetic SN. Of course, the large explosion energy claimed for SN 1998bw is based on the assumption was spherically symmetric. There is growing evidence that SN Ib/c show a small polarization, implying that their explosions might be somewhat aspherical.

Other evidence linking Type Ib/c SN to Wolf-Rayet precursors includes pre-SN imaging of M74, the host galaxy of SN 2002ap (Type Ic SN), which led Smartt et al. (2002) to conclude that a Wolf-Rayet star was a viable candidate for the immediate progenitor..

10.2 Hypernovae and Gamma Ray Bursts

It is only within the past few years that Gamma Ray Bursts (GRBs), in which $0.1\text{--}1$ MeV photons are detected for a few seconds, have been shown to be of cosmological origin, typically in the range $z=0.5$ to 2 . There are two, potentially distinct populations of GRBs, peaking at 0.2s and 20s (Fishman & Meegan 1995), which may have different origins, the latter being most common.

The rapid (millisecond) γ -ray variability of GRBs implies a compact object, of size smaller than a light millisecond (10^7 cm) and mass not greater than $\sim 30 M_{\odot}$. The total energy released in γ -rays corresponds to a rest mass energy of a fraction of a solar mass. The most natural way for triggering a GRB event is the accretion of a fraction of a solar mass onto a black hole. Therefore, most cosmological GRB models have at their basis gravitational collapse of a (several) solar mass progenitor to a black hole. The duration of long bursts implies that some fraction of the progenitor mass accretes via a disk. The so-called 'collapsar' model for long duration bursts, favours a rapidly rotating (massive) stellar core which collapses to form a rotating black hole plus an accretion torus (Woosley et al. 1999).

The first direct link between core collapse SNB and GRBs was established by the 'hypernova' SN 1998bw which was spatially coincident with GRB 980425 (Galama et al. 1998). The host galaxy for this source was established to lie at $z=0.0085$, such that the GRB was found to be intrinsically many orders of magnitude fainter than classical GRBs (Woosley et al. 1999). Consequently, the link

between ‘hypernovae’ and GRBs still awaited more general confirmation. Indeed, there are some hypernovae for which no GRB counterpart has been proposed, including SN 2002ap which was less energetic than SN 1998bw. It is possible that weaker explosions are less efficient in collimating the γ -rays to give rise to a GRB, or there is an inclination of the beam axis with respect to the line-of-sight to the SN.

Subsequently, Reeves et al. (2002) observed the X-ray afterglow of GRB 011211 using XMM-Newton, revealing spectral lines exhibiting an outflow velocity of $0.08c$, plus evidence of enriched α -group elements, characteristics unique to core collapse SN. However the GRB was too faint for the SN to be observed optically.

Very recently, the extremely bright, long duration GRB 030329 (Stanek et al. 2003), at a low redshift of $z=0.1685$, has been found to reveal a ‘hypernova’ spectrum SN 2003dh. Other cases in which supernovae has been inferred from light curves and colours of GRB afterglows, though this represented direct proof that a subset of classical GRBs originate from SN. From the accumulated evidence at hand, it is clear that there is a direct link between SN originating from core collapse massive stars and long duration GRBs.

11 Further Reading

Current research relating to Wolf-Rayet stars can be found in the proceedings of IAU Symposium 212 (van der Hucht et al. 2003), whilst Luminous Blue Variables are discussed in Nota & Lamers (1997). A recent volume in the Lecture Notes in Physics series discusses Supernova and GRBs (Weiler 2003).

References

- Barba R.H., Niemela V.S., Baume G. & Vazquez R.A., 1995, ApJ 446, L23
- Barlow M.J., Hummer D.G., 1982, in Proc IAU Symp 99, *Wolf-Rayet Stars: Observations, Physics, Evolution*, C. de Loore & A. Willis, eds., (Dordrecht: Reidel), p 387
- Beals C.S. & Plaskett H.H., 1935, Trans. IAU, 5, 134
- Blöcker, T., Balega, Y. & Hofmann, K.-H. et al. 1999, A&A 348, 805
- Blum R.D., Schaerer D., Pasquali A., Heydari-Malayeri M., Conti P.S. & Schmutz W., 2001, AJ 122, 1875
- Bohannon B. & Crowther P.A., 1999 ApJ 511, 374
- Cappellaro E., Evans R. & Turatto M., 1999, A&A 351, 459
- Chapman J.M., Leitherer C., Koribalski B., Bouter R. & Storey M., 1999, ApJ 518, 890
- Clark J.S. & Negueruela I., 2002, A&A 396, L25
- Conti P.S., 1976, in Proc IAU Symp 70, *Be and shell stars*, Slettebak, A. ed., (Dordrecht: Reidel), p. 447
- Conti P.S., 1984, in Proc IAU Symp 105, *Observational Tests of the Stellar Evolution Theory* Maeder A., & Renzini A., eds. (Dordrecht: Reidel), p.233
- Conti P.S., 1997, in *Luminous Blue Variables: Massive Stars in Transition*, Nota A. & Lamers H.J.G.L.M., eds, (San Francisco: ASP), ASP Conf. Ser 120, p.161

- Conti P.S., Leep M.E. & Perry D.N., 1983, ApJ 268, 228
- Conti P.S. & Massey P., 1989, ApJ 337, 251
- Crowther P.A., 1997, in *Luminous Blue Variables: Massive Stars in Transition*, Nota A. & Lamers H.J.G.L.M., eds, (San Francisco: ASP), ASP Conf. Ser 120, p.51
- Crowther P.A., 2000, A&A 356, 191
- Crowther P.A., Hillier D.J. & Smith L.J., 1995a, A&A 293, 172
- Crowther P.A., Hillier D.J. & Smith L.J., 1995b, A&A 293, 403
- Crowther P.A., Smith L.J. & Hillier D.J., 1995c, A&A 302, 457
- Crowther P.A., Smith L.J. & Willis A.J., 1995d, A&A 304, 269
- Crowther P.A. & Smith L.J., 1997, A&A 320, 500
- Crowther P.A. & Dessart L., 1998, MNRAS 296, 622
- Crowther P.A., De Marco O. & Barlow M.J., 1998, MNRAS 296, 367
- Crowther P.A., Fullerton A.W., Hillier D.J. et al., 2000, ApJ 538, L51
- Crowther P.A., Dessart L., Hillier D.J. et al. 2002a, A&A 392, 653
- Crowther P.A., Hillier D.J., Evans C.J. et al. 2002b, ApJ 579, 774
- Crowther P.A., Drissen L., Abbott J.B., Royert P. & Smartt S.J., 2003, A&A in press (astro-ph/0302475)
- Damineli A., Conti P.S. & Lopes D.F., 1997, New Astron, 2, 107
- Damineli A., Kaufer A., Wolf B. et al. 2000, ApJ 528, L101
- de Jager, C., 1998, A&A Rev 8, 145
- de Mello D.F., Schaerer D., Heldmann J. & Leitherer C., 1998, ApJ 507, 199
- de Koter A., Hubeny I., Heap S.R. & Lanz T., 1994, ApJ 435, L71
- de Koter A., Heap S.R. & Hubeny I., 1997, ApJ 477, 792
- De Marco O., Schmutz W., Crowther P.A. et al. 2000, A&A 358, 187
- Dessart L. & Owocki S.P., 2002, A&A 383, 1113
- Dessart L., Crowther P.A., Hillier D.J., et al. 2000, MNRAS 315, 407
- Dray L.M., Tout C.A., Karakas A.I. & Lattanzio J., 2003, MNRAS 338, 973
- Drissen L., Crowther P.A., Smith L.J. et al. 2001, ApJ 546, 484
- Eenens P.R.J. & Williams P.M., 1994, MNRAS 269, 1082
- Esteban C., Vilchez J.M., Smith L.J. & Clegg R.E.S., 1992, A&A 259, 629
- Figer D., McLean I.S. & Najarro F., 1997, ApJ 486, 420
- Figer D., Najarro F., Morris M. et al. 1998, ApJ 506, 384
- Figer D., McLean I.S. & Morris M., 1999a, ApJ 514, 202
- Figer D., Morris M., Geballe T.R. et al. 1999b, ApJ 525, 759
- Fishman G.J. & Meegan, C.A., 1995, ARA&A 33, 415
- Galama T.J., Vreeswijk P.M., van Paradijs J. et al. 1998, Nat 395, 670
- Gräfener G., Koesterke L. & Hamann W.-R., 2002, A&A 387, 244
- Grosdidier, Y., Moffat, A.F.J., Joncas, G. & Acker, A., 1998, ApJ 506, L127
- Gummersbach, C.A., Zickgraf F.-J. & Wolf B., 1995, A&A 302, 409
- Hamann W.-R. & Koesterke L., 1998, A&A 335, 1003
- Harries T.J., Hillier D.J. & Howarth I.D., 1998, MNRAS 296, 1072
- Heap S.R., Altner B., Ebbets D. et al. 1991, ApJ 377, L29

- Herald J.E., Hillier D.J. & Schulte-Ladbeck R.E., 2001, ApJ 548, 932
- Hillier D.J., 1991, A&A 247, 455
- Hillier D.J. & Miller D.L. 1998, ApJ 496, 407
- Hillier D.J., Crowther P.A., Najarro F. & Fullerton A.W., 1998, A&A 340, 483
- Hillier D.J., Davidson K., Ishibashi K. & Gull T., 2001, ApJ 553, 837
- Homeier N.L., Blum R.D., Conti P.S. & Damineli A., 2003, A&A 397, 585
- Howarth I.D. & Schmutz W., 1992, A&A 261, 503
- Humphreys R.M. & Davidson K., 1979, ApJ 232, 409
- Humphreys R.M., Davidson K. & Smith N., 2002, AJ 124, 1026
- Iwamoto, K. et al. 1998, Nat 395, 672
- Iwamoto, K., Nakamura, T., Nomoto, K. et al. 2000, ApJ 534, 660
- Jones, T.J., Humphreys, R.M., Gehrz, R.D. et al. 1993, ApJ 411, 323
- Josselin, E., Blommaert J.A.D.L., Groenewegen, M.A.T., Omont, A. & Li, F.L., 2000, A&A 357, 225
- Jura, M. & Kleinmann, S.G., 1990, ApJS 73, 769
- Koesterke L. & Hamann W.-R., 1995, A&A 299, 503
- Kingsburgh R.L., Barlow M.J. & Storey P.J., 1995, A&A 295, 75
- Lamers H.J.G.L.M., Zickgraf F.-J., de Winter D., Houziaux L. & Zorec J., 1998, A&A 340, 117
- Lamers H.J.G.L.M., Nota A., Panagia N., Smith L.J. & Langer N., 2001, ApJ 551, 764
- Langer N., Garcia-Segura G. & MacLow M.-M., 1999, ApJ 520, L49
- Leitherer C., 1997, in *Luminous Blue Variables: Massive Stars in Transition*, Nota A. & Lamers, H.J.G.L.M., eds, (San Francisco: ASP), ASP Conf. Ser 120, p.58
- Leitherer C., Schmutz W., Abbott D.C., Hamann W.-R. & Wessolowski U., 1989, ApJ 346, 919
- Leitherer C., Robert C. & Drissen L., 1992, ApJ 401, 596
- Leitherer C., Chapman J.M. & Koribalski B., 1997, ApJ 481 898
- Lennon D.J., Kudritzki R.-P., Haser S., Fitzpatrick E.L. & Walborn N.R., 1997, in *Luminous Blue Variables: Massive Stars in Transition*, Nota A. & Lamers H.J.G.L.M., (eds), (San Francisco: ASP), ASP Conf. Ser 120, p.90
- Lepine S., Moffat A.F.J., St-Louis N. et al. 2000, AJ 120, 3201
- Maeder A. & Meynet G., 1994, A&A 287, 803
- Martins F., Schaerer D. & Hillier D.J., 2002, A&A 382, 999
- Massey P., 1996, in: Proc. 33rd Liège Intl. Astrophysics Coll., *Wolf-Rayet stars in the framework of stellar evolution*, p.361
- Massey P. & Johnson O., 1998, ApJ 505, 793
- Massey P. & Duffy A.S., 2001, ApJ 550, 713
- Massey P. & Holmes S., 2002, ApJ 580 L35
- Massey P., Armandroff T.E. & Conti P.S., 1992, AJ 103, 1159
- Massey P., DeGioia-Eastwood K. & Waterhouse E., 2001, AJ 121, 1050
- Moore B.D., Hester J.J. & Scowen P.A., 2000, AJ 119, 2991
- Morse J.A., Davidson K., Bally J., Ebbets D., Balick B. & Frank A., 1998, AJ 116, 2443
- Najarro F., Krabbe A., Genzel R. et al. 1997a, A&A 325, 700

- Najarro F., Hillier D.J. & Stahl O. 1997b, A&A 326, 1117
- Nota A. & Lamers, H.J.G.L.M. (eds), *Luminous Blue Variables: Massive Stars in Transition*, ASP Conf. Ser 120, (San Francisco, ASP)
- Nugis T. & Lamers H.J.G.L.M., 2000, A&A 360, 227
- Prinja R.K., Barlow M.J. & Howarth I.D., 1990, ApJ 361, 607
- Puls, J., et al. 1996, A&A, 305, 171
- Reeves, J.N. et al. 2002, Nat 416, 512
- Reid, N., Tinney, C. & Mould, J., 1990, ApJ 348, 98-119
- Royer P., Smartt S.J., Manfroid J. & Vreux J.-M., 2001, A&A 366, L1
- Sanduleak, N., 1977. *Inf. Bull. Var. Stars.*, 1304
- Schaerer D. & Maeder A., 1992, A&A 263, 129
- Schild H., Crowther P.A., Abbott J.B. & Schmutz W., 2003, A&A 397, 859
- Schweickhardt J., Schmutz W., Stahl O., Szeifert Th. & Wolf B., 1999, A&A 347, 127
- Shapley A.E., Steidel C.C., Pettini M. & Adelberger K.L., 2003, ApJ 588, 65
- Smartt S.J., Vreeswijk P.M., Ramirez-Ruiz E. et al. 2002, ApJ 572, L147
- Smith L.F. 1968a, MNRAS 138, 109
- Smith L.F., 1968b, MNRAS 140, 409
- Smith L.F. & Hummer D.G., 1988, MNRAS 230, 511
- Smith L.F., Shara M.M. & Moffat A.F.J., 1990, ApJ 358, 229
- Smith L.F., Maeder A. & 1991, A&A 241, 77
- Smith L.F., Shara M.M. & Moffat A.F.J., 1996, MNRAS 281, 263
- Smith L.J., Crowther P.A. & Prinja R.K., 1994, MNRAS 281
- Smith L.J., Nota A., Pasquali A. et al. 1998, ApJ 503, 278
- Smith L.J., Norris R.P.F. & Crowther P.A., 2002, MNRAS 337, 1309
- St-Louis N., Moffat A.F.J., Turbide L. & Bertrand J.-F., 1998, in *Boulder-Munich II: Properties of Hot, Luminous Stars*, Howarth I.D. ed., (San Francisco: ASP), ASP Conf. Ser 131, p.326
- Stanek, K.Z., Knapp, G.R., Young, K. & Phillips, R.G. 1995, ApJS 100, 169
- Stanek K.Z., Matheson T., Garnavich P.M. et al. 2003, (astro-ph/0304173)
- Sylvester, R., Skinner, C.J. & Barlow, M.J. 1998, MNRAS 301, 1083
- Tubbesing S., Kaufer A., Stahl O. et al. 2002, A&A 389, 931
- van der Hucht K.A., 2001, New Astronomy Rev, 45, 135
- van der Hucht K.A., Herrero A. & Esteban C. (eds), 2003, IAU Symposium 212: *A Massive Star Odyssey, from Main Sequence to Supernova*, (San Francisco: ASP)
- van Loon, J.Th., Groenewegen, M.A.T., de Koter, A. et al. 1999, A&A 351, 559
- Vink J.S., de Koter A. & Lamers H.J.G.L.M., 2001, A&A 369, 574
- Walborn N.R., 1973, ApJ 179, 517
- Walborn N. R. 1977, ApJ 215, 53
- Walborn N.R., 1982, ApJ, 256, 452
- Walborn N.R., 1986, in Proc. IAU Symp. 116, *Luminous Stars and Associations in Galaxies* De Loore, C.W.H., Willis A.J. & Laskarides P., eds, (Dordrecht: Reidel), p. 185
- Walborn N.R., Howarth I.D., Lennon D.J. et al., 2002, AJ 123, 2754

- Weiler K. (ed), *Supernovae and Gamma-Ray Bursts*, Lecture Notes in Physics, 598,
(Berlin: Springer-Verlag)
- Willis A.J., van der Hucht K.A., Conti P.S. & Garmany C.D., 1986, A&AS 63, 417
- Woosley S.E., Eastman, R.G. & Schmidt, B.P., 1999, ApJ 516, 788
- Wright A.E., Barlow M.J., 1975, MNRAS 170, 41
- Zickgraf F.-J., Wolf B. Stahl O. et al. 1985, A&A 143, 421
- Zickgraf F.-J., Kovacs J. Wolf B. et al. 1996, A&A 309, 505